PATENT APPLICATION

Docket Number: 037003-0280705 Client Reference: 1995-30-0233CP3

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In Re the Application of ANDERSON et al.

Group Art Unit: 1644

Application No.: 10/073,138

Examiner: Phillip Gambel

Filed: February 13, 2002

Confirmation No.: 7969

For: IDENTIFICATION OF UNIQUE BINDING INTERACTIONS BETWEEN CERTAIN ANTIBODIES AND THE HUMAN B7.1 AND B7.2 CO-STIMULATORY ANTIGENS

INFORMATION DISCLOSURE STATEMENT

Commissioner for Patents P. O. Box 1450 Alexandria, VA 22313-1450

Sic:

Pursuant to 37 CFR 1.56, the attention of the Patent and Trademark Office is hereby directed to the reference(s) listed on the attached PTO-1449. The references were cited by the applicant or examiner during the prosecution of the parent application, U.S. Application No. 09/383,916, filed on August 26, 1999. Thus, copies of the references listed in the attached PTO-1449 are not attached. It is respectfully requested that the information be expressly considered during the prosecution of this application, and that the reference(s) be made of record therein and appear among the "References Cited" on any patent to issue therefrom.

This Information Disclosure Statement is being filed before the mailing date of the first official action on the merits in the present application. No certification or fee is required.

Respectfully submitted,

RACarl

Thomas A. Cawley, Jr., Ph.D. Registration Number 40,944

Date: February 6, 2004 PILLSBURY WINTHROP LLP Telephone: (703) 905-2000 Facsimile: (703) 905-2500

P.O. Box 10500 McLean. VA 22102 TAC/JBM:ntb

	15:24	From-PILLSBURY WINTHROP +703-905-2500 T-868 P.005/006 F-500
AAR	Dautig	ny, et al., "Molecular cloning and nucleotide sequence of a cDNA clone coding for rat brain myelin hippid," FEBS Lett., 1985, 188(1):33-36.
BBR	Delab	ie, et al. "The B7/BB1 antigen is expressed by Reed-Sternberg Colls of Hospital (1993, 82:2845-52.
CCR		
DDR	Durie	, et al., "The role of CD40 and its ligand (gp39) in peripheral and central district
EER	Durie	, et al , "Prevention of collagen-induced artificis with all artificody to good, are injuried.
FFR	Falin	i, et al., "Response of refractory Hodgkin's disease to monocional anti-CD30 immultidoxiii," Earness,
GGR	Free	man, et al., "Uncovering of functional alternative CTLA-4 counter-receptor in 57-deficient times,"
ннR	Free	man, et al., "Cloning of B7-2: a CTLA-4 counter-receptor that costimulates numan if cell profileration,
IIR	Gerr	itse, et al., "CD40-CD40 ligand interactions in experimental allergic encephalothyeitis and multiple
JJR	Gott	tieb, et al., "Results of a single-dose dose-escalating trial of an anti-B7.1 antibody (IDEC-1147 in putients
KKR	Gott	heb, et al., "Clinical and histologic response to single-dose treatment of moderate to severe positions to single-dose treatment of moderate to severe positions." J. Am. Acad. Dermatol., 2002, 47:693-700.
LLR	Guil	nan, et al., "Pivotal role of the B7:CD28 pathway in transplantation tolerance and tumor intimating," was a
ммі	R Haf	er, et al., "The potential of restricted T cell recognition of myelin basis protein epitopes in the therapy of
NNF	R Har	inaran, et al. "Therapeutic activity of IDEC-114 (anti-CD80) and muximab (Kituxane) in B-carrymphoma
00	R Hat	hoock, et al., "Identification of an alternative CTLA-4 ligand costimulatory for 1 cell activation," Science,
PP	CD	3, 282:905-907. lenbaugh, et al., "The human T cell antigen gp39, a member of the TNF gene family, is a ligand for the 40 receptor expression of a soluble form of gp39 with B cell co-stimulatory activity," EMBO J., 1992, 12) 4313-4321
QQ	0 1/2	TIME LINE SUPPRESSIVE Therapy " Curr Opin Immunol., 1992, 4:553-560.
RR	R Ka	pus, et al. "CD4+ suppressor cells differentially affect the production of IFN-y by effector cells differentially affect the IFN-y by effector cel
ss		*Characters based strategies for drug design and discovery," Science, 1992, 257, 1074-d2.
11	R La	man, et al., "The role of gp39 (CD40 ligand) in EAE and MS," Journal of NeuroInfinitiology, 1334, 54(1
- Juu	R Le	derman, et al. "Identification of a novel surface protein on activated CD4+ 1 cells that induces comact-
V	R Lic	er, et al., "Suppression of experimental autoimmune encephalomyellus by oral authinistation of myoming
W'	WR Li	isley, et al., "The role of the CD28 receptor during T cell responses to antigen," Ann. Rev. Intitution, 133.
xx	(R Li	i. et al., "Blockade of CD28/CTLA-4-B7 co-stimulatory pathway in colitic SCID mice." Digestive Disease
~	R M	Cafferty, et al ," Phage antibodies. filamentous phage displaying antibody variable domains, mature, 133
Z	R M	iller, et al., "Antigen-driven bystander suppression after oral administration of antigens, "J EAP. Med., 132
1	AAR M	okhtarion, et al., "Adoptive transfer of myelin basic protein-sensitized 1 cells produces chiofic relapsing
1 1	BBR M	orrison, et al., "Chimeno human antibody molecules, mouse antigen-binding domains with human constant
1 1	CCR N	gion domains. Froc. Near Actas, Set 3331, and its grant process of antigen-presenting cells a number of the state of the s
1		

Fab-08-2004 15:24 From-PILLSBURY WINTHROP +703-805-2500 T-868 P.006/006 F-500

	Nickoloff, et al., "T lymphocytes in skin lesions of psoriasis and mycosis fungoides express B7-1; a ligand for CD28," <i>Blood</i> , 1994, 83:2580-2586.
FFFR	Noelle, et al., "A 39-kDa protein on activated helper T cells binds CD40 and transduces the signal for cognities and transduces the signal for cognities are represented by the signal for cognities and transduces the signal for cognities are represented by the signal for cognities are represented by the signal for cognities are represented by the signal for cognities and transduces the signal for cognities are represented by the signal for cognities are repre
GGGR	Olsson, et al., "Human-human monoclonal antibody-producing hybridomas, technical aspects," <i>Meth.</i>
HHHR	Pernn, et al., "Opposing effects of CTLA4-ig and anti-CD80 (B7-1) plus anti-CD86 (B7-2) on experimental allocological properties." J. Neuroimmunol., 1996, 65:31-39.
IIIR	Pesoa, et al., Regulation of experimental allergic encephalomyelitis. Part 5. Role of the recepient in
אררי	Pettinelli, et al., "Adoptive transfer of experimental allergic encephalomyelitis in SJDJ mice after in Vibulativation of lymph node cells by myelin basic protein: requirement for Lyt 1" 2" T lymphocytes," J. Immunol., 1979, 1977, 1979, 1977, 1979, 1977, 1979, 1977, 1979, 197
3	Skolnick, et al., "From genes to protein structure and function: novel applications of computational Skolnick, et al., "From genes to protein structure and function: novel applications of computational Skolnick, et al., "From genes to protein structure and function: novel applications of computational Skolnick, et al., "From genes to protein structure and function: novel applications of computational Skolnick, et al., "From genes to protein structure and function: novel applications of computational Skolnick, et al., "From genes to protein structure and function: novel applications of computational Skolnick, et al., "From genes to protein structure and function: novel applications of computational Skolnick, et al., "Trends in Biotechnology, 2000, 18:34-9
1	Sobel, et al. "Acute experimental allergic encephalomyelitis in SJL/J mice induced by a synthetic peptide of
	Stamenkovic, et al., "A B-lymphocyte activation molecule related to the nerve growth factor receptor and increase by cytokines in carcinomas." EMBO J., 1989, 8(5),1403-1410.
L.	Suvas, et al., "Distinct role of CD80 and CD86 in the regulation of the activation of B cell and B cell
1	Takeda, et al. "Construction of chimaeric processed immunoglobulin genes containing mouse variable and
	Teng, et al., "Construction and testing of mouse-human heteromyelomas for numan monocional antibody
1	Tuohy, et al., "Identification of an encephalitogenic determinant of myelin proteolipid protein for SJL mice, 3
I	Van der Veen, et al., "The adoptive transfer of chronic relapsing experimental altergic encephalotryellus with
1	Ward, et al., "Binding activities of a repertoire of single immunoglobulin variable domains secretary from
	Ward, et al., "Blocking of adhesion molecules in vivo as anti-inflammatory therapy," Ther Immunoi., 1994,
UUUI	R Yi-qun, et al. "Differential requirements for co-stimulatory signals from B7 family members by resting versus recently activated memory T cells towards soluble recall antigens." <i>Int. Immunol.</i> , 1996, 8 37-44.
xaminer	Date Considered: